

# Circles

4<sup>TH</sup> AUTUMN SERIES

MODEL SOLUTIONS


## Problem 1.

Kua would like to draw 7 circles on a piece of paper so that every two circles intersect at two distinct points. And he would love it if there were exactly 7 points, in which 2 or more circles intersect. Show him how this is possible.

(Filip Bialas)

SOLUTION:

We can draw an equilateral triangle with its centre and midpoints of its sides. And then draw circles as in the diagram.



More precisely, denote the vertices of the triangle by  $A, B, C$ , midpoints of the opposite sides by  $D, E, F$  and its centre by  $G$ . Then it suffices to consider circles with diameters  $AG, BG, CG, AB, BC, AC$  and circumcircle of  $DEF$ .

POZNÁMKY:

Většina řešitelů přišla s podobným obrázkem jako ve vzoráku. Pouze dva řešitelé našli mnohem méně symetrický obrázek, kde jedním z vrcholů procházelo hned šest kružnic. U všech ostatních, kteří tuhle úlohu zdárně vyřešili, šlo o kombinatoricky stejnou konstrukci – průsečíky šlo vždy pojmenovat tak, aby na nakreslených sedmi kružnicích ležely stejně pojmenované body jako na těch ve vzorovém řešení. I když u některých vypadal obrázek na první pohled o dost jinak.

*Matoušovi Šafránkovi* jsem udělil +i za to, že si uvědomil, že při nalézání obrázku nemusí používat pouze kružnice, ale také přímky. Vzniklý obrázek pak již jen stačí zinvertovat podle libovolné kružnice se středem v bodě, který neleží na žádné z nakreslených přímek a kružnic. Pokud poslední větě nerozumíte, ale chtěli byste jí rozumět, stačí se podívat do druhého dílu letošního seriálu. (Filip Bialas)

**Problem 2.**

Let  $\omega_1, \omega_2$  be two circles with centres  $O_1, O_2$  intersecting at points  $X, Y$  such that  $\angle O_1XO_2 = 90^\circ$ . Let  $D$  be the intersection of  $O_1O_2$  and  $\omega_1$  such that  $O_1$  lies between  $D$  and  $O_2$ . Let  $P$  be the intersection of  $DX$  and  $\omega_2$  distinct from  $X$ . Prove that  $PO_2$  is perpendicular to  $O_1O_2$ .

(Pavel Hudec)

SOLUTION:

Since  $DO_1$  and  $XO_1$  are radii of  $\omega_1$ , triangle  $DO_1X$  is isosceles. Using a similar argument, we get that  $PO_2 = XO_2$ , so triangle  $PO_1X$  is also isosceles. Hence we have

$$\angle XDO_1 = \angle DXO_1,$$

$$\angle XPO_2 = \angle PXO_2.$$

The points  $D, X, P$  are colinear, therefore


$$\angle DXO_1 + \angle O_1XO_2 + \angle O_2XP = 180^\circ,$$

$$\angle DXO_1 + \angle O_2XP = 90^\circ,$$

$$\angle XDO_1 + \angle O_2PX = 90^\circ.$$

Because the sum of internal angles in the triangle  $DPO_2$  is  $180^\circ$ , we can get the desired conclusion:

$$\angle DO_2P = 180^\circ - \angle O_2DX - \angle O_2PX = 90^\circ.$$



POZNÁMKY:

Úlohu vyřešila většina řešitelů správně. Jen připomínám, že řešení má být pochopitelné i bez příloženého obrázku, tedy je potřeba v textu definovat i všechny úhly, které nějak označíme, a provést všechny výpočty, ne se pouze odkázat na obrázek. („madam Verča“ Hladíková)

### Problem 3.

Jáchym drew three circles on a whiteboard. The circles had radii 2, 3, and 3 and each two were externally tangent. Then he drew the circle  $\omega$  that is internally tangent to all three of them. Help him calculate the radius of  $\omega$ .

(Lucien Šíma)


SOLUTION:

Let  $\omega_1, \omega_2, \omega_3$  be the three circles and  $O_1, O_2, O_3$  their respective centres ( $\omega_1$  has a radius of 2). Let  $M$  be the midpoint of  $O_2O_3$ . The configuration is symmetric with respect to the common tangent line of  $\omega_2$  and  $\omega_3$ , namely the line  $O_1M$ , therefore  $O_1M$  is perpendicular to  $O_2O_3$  and the centre of  $\omega$  lies on  $O_1M$ . The length of  $O_1M$  can be obtained from the Pythagorean theorem as

$$O_1M = \sqrt{O_1O_2^2 - O_2M^2} = \sqrt{5^2 - 3^2} = 4.$$

Let  $X$  be the intersection of  $O_1M$  with  $\omega_1$  that is further away from  $M$  and let  $Y$  and  $Z$  be the intersections of  $O_2O_3$  with  $\omega_2$  and  $\omega_3$  respectively distinct from  $M$ . Then the centre of  $\omega_1$  lies on the line  $XM$  and  $XM = XO_1 + O_1M = 2 + 4$ . Similarly,  $O_2$  and  $O_3$  lie on the line  $MY$  (and  $MZ$ ) and  $MY = MZ = 3 + 3 = 6$ . Therefore if we draw a circle with the centre  $M$  and radius 6, it will be internally tangent to all three circles at points  $X, Y$ , and  $Z$ .

The radius of  $\omega$  is 6.



POZNÁMKY:

Většina si s úlohou hravě poradila, ale několik řešitelů místo poloměru kružnice, která má se zadanými kružnicemi vnitřní dotyk (*internally tangent*), spočítalo poloměr kružnice, která má dotyk vnější (*externally tangent*), a tím si vysloužili nula bodů. (Hedvika Ranošová)

**Problem 4.**

Let  $ABCD$  be a cyclic quadrilateral with circumcentre  $O$  such that  $AC$  and  $BD$  are perpendicular. Let  $\omega_1, \omega_2, \omega_3,$  and  $\omega_4$  be circles, where the diameters of these circles are  $AO, BO, CO,$  and  $DO$  respectively. Finally, let  $P, Q, R,$  and  $S$  be the intersections of  $\omega_1$  with  $\omega_2, \omega_2$  with  $\omega_3, \omega_3$  with  $\omega_4,$  and  $\omega_4$  with  $\omega_1$  respectively, distinct from  $O$ . Prove that  $PQRS$  is a rectangle.

(Jáchym Solecký)

SOLUTION:

First, we would like to prove that  $P, Q, R, S$  are the midpoints of segments  $AB, BC, CD, DA$  respectively. Have a look at point  $P$  which lies on two circles with diameters  $AO$  and  $BO$ . Therefore, from Thales' theorem, we know that the angles  $APO$  and  $BPO$  are right. So the size of the angle  $APB$  is

$$\angle APB = \angle AOP + \angle BOP = 180^\circ.$$

This means that  $P$  lies on the segment  $AB$ . In addition, triangle  $AOB$  is isosceles with the base  $AB$  (because  $AO$  and  $BO$  are radii of the circumcircle of the quadrilateral  $ABCD$ ). So the altitude from point  $O$  to the base is also a median and therefore  $P$  is the midpoint of  $AB$ . By analogy, we get the same result for the rest of the points  $Q, R, S$ .

As a result of that, we know that the lines  $PQ, QR, RS, SP$  are the midlines of the triangles  $ABC, BCD, CDA, DAB$  respectively.

Then, as we know, in triangle  $KLM$  the midline connecting the midpoints of  $KL$  and  $KM$  is parallel to the third side  $LM$ . This implies that  $PQ$  and  $SR$  are parallel to their base  $AC$ , analogically  $QR$  and  $SP$  are parallel to their base  $BD$ . Moreover,  $AC$  is perpendicular to  $BD$  and this all together means that  $PQ$  and  $SR$  are perpendicular to  $QR$  and  $SP$ , so  $PQRS$  is a rectangle.

POZNÁMKY:

Většina došlých řešení dostala plný počet bodů. Někteří řešitelé považovali některé věci za zřejmé, třeba že body  $P, Q, R, S$  leží na úsečkách  $AB, BC, CD, DA$ , ačkoli by zasloužily důkaz. Proto jim byl občas strhnut bod. Jinak bylo všechno v pořádku a neměl jsem žádné velké výtky.

(Fila Čermák)

**Problem 5.**

Let  $\omega_1$  and  $\omega_2$  be two circles externally tangent at  $T$ . Let  $C$  be a point on  $\omega_2$  such that the tangent at  $C$  intersects  $\omega_1$  at two distinct points  $X$  and  $Y$ . Now define  $P$  as the intersection of  $CT$  and  $\omega_1$  distinct from  $T$ . Show that  $PXY$  is an isosceles triangle.

(Pavel Hudec)

SOLUTION:

Let  $O_1, O_2$  be the centres of  $\omega_1, \omega_2$ . Consider the homothety with center  $T$  which sends  $\omega_1$  to  $\omega_2$ . The image of the line  $O_1P$  is  $O_2C$ , therefore  $O_1P \parallel O_2C$  (since in a homothety, every line is parallel to its image). It follows that  $O_1P \perp XY$  because  $XY$  is tangent to  $\omega_2$  and perpendicular to  $O_2C$ . Finally,  $O_1$  lies on the perpendicular bisector of  $XY$ , hence  $O_1P$  and the perpendicular bisector coincide. We have proven that  $P$  lies on the perpendicular bisector of  $XY$  which implies  $PX = PY$  and so  $PXY$  is isosceles.

POZNÁMKY:

Téměř všechna řešení byla správná a většina z nich postupovala podobně jako vzorové řešení. Ukázalo se, že úlohu je možné řešit mnoha různými způsoby. Část řešitelů například využívala průsečíku  $XY$  a společné tečny ke kružnicím v bodě  $T$ .

(Josef Minařík)

### Problem 6.

There are  $2n$  points on a circle labelled  $1, 2, \dots, 2n$  in some order. We define a pairing as a set of  $n$  segments between these points such that every point is an endpoint of exactly one of the segments. For a segment connecting points labelled  $a$  and  $b$ , we say its value is the number  $|a - b|$ . Finally, we say a pairing is good, if the sum of values of all  $n$  segments is equal to  $n^2$ . Show that for any initial order of labels there exists a good pairing such that no two segments intersect.

(Pavel Hudec)

SOLUTION:

First, colour points labelled  $1, 2, \dots, n$  as red and points labelled  $n+1, n+2, \dots, 2n$  as blue. We will say a pairing is *codenamesish* if every segment connects a red point with a blue point and no two segments intersect. Assume that we have a codenamesish pairing. Then any blue point (labelled by  $a$ ) would contribute to the sum of values of all  $n$  segments by  $a$ , whereas any red point (labelled by  $b$ ) would contribute by  $-b$ . Therefore the sum of values is equal to

$$\left( (n+1) + (n+2) + \dots + 2n \right) - (1 + 2 + \dots + n) = n^2.$$


We will present two proofs of existence of a codenamesish pairing.

FIRST SOLUTION (CONNECTING NEIGHBORING POINTS):

Use induction on  $n$ . The base case  $n = 1$  is clear. Now suppose there are  $n$  blue and  $n$  red points on the circle. Note that there exist two neighboring points of opposite colours. Connect them by a new segment. By the induction hypothesis there exists a codenamesish pairing of the remaining points. These all lie in one half-plane defined by the new segment, therefore no other segment can intersect the new segment. We can then conclude that the constructed pairing is indeed codenamesish.

SECOND SOLUTION (REDUCING THE NUMBER OF INTERSECTIONS):

Consider a pairing where every segment connects a red point with a blue point with the smallest number of intersecting pairs of segments. For the sake of contradiction, suppose that there exist two intersecting segments  $AC$  and  $BD$ . Now, connect the points  $A, B, C, D$  in the other way such that both segments still connect a red point with a blue point. WLOG let the connected pairs of points now be  $AB$  and  $CD$ .



Notice that  $AB$  and  $CD$  don't intersect, which leaves us with one fewer intersection points. If any other segment intersects both  $AB$  and  $CD$ , then it has to intersect both  $AC$  and  $BD$ , as the quadrilateral  $ABCD$  is convex. If it intersects just one of them, say  $AB$ , it has to intersect at least one of the diagonals as well, since  $AB$  and parts of  $AC$  and  $BD$  form a triangle. Therefore the number of intersections after the swap decreased by at least one. This contradicts the minimality property of the pairing.

POZNÁMKY:


Úloha dopadla velmi dobře, pouze několik málo řešitelů nad ní zaváhalo. Při konstrukci hledaného párování řešitelé využívali množství různých přístupů. Většina řešení byla podobná prvnímu vzorovému, ovšem byla k vidění i řešení používající diskrétní spojitost nebo některý z možných extrémálních principů. (Pavel Hudec)

**Problem 7.**

Let  $ABCD$  be a parallelogram such that  $\angle DAB$  is obtuse. Then, let  $M$  be the midpoint of  $AB$  and  $E$  be the intersection of the circumcircle of  $DAB$  and the line  $DM$  distinct from  $D$ . Finally, let  $H$  be the point on  $DA$  such that  $\angle AHB = 90^\circ$ . Prove that  $C, D, H,$  and  $E$  are concyclic. (Matěj Doležálek)

SOLUTION:

Let  $\omega$  be the circumcircle of  $DAB$  and let  $K$  be the point on  $BC$  such that  $\angle AKB = 90^\circ$ . Then  $AHBK$  is a rectangle since  $AH$  and  $BK$  are parallel and two opposite angles are right.



We shall prove that  $D, C, K, H$  are concyclic and that  $D, K, E, H$  are concyclic. That will mean that all four points  $C, D, H, E$  lie on the circumcircle of  $DKH$ , solving then the problem.

First, the diagonals of a rectangle have the same length, so  $HK = AB$ . Since  $DH$  and  $CK$  are parallel, this means that  $DCKH$  is an isosceles trapezoid and thus cyclic. Second, because the diagonals of a rectangle halve each other,  $M$  is the midpoint of both  $AB$  and  $HK$ , so the power<sup>1</sup> of  $M$  with respect to  $\omega$  gives us

$$DM \cdot ME = AM \cdot MB = HM \cdot MK.$$

This means that  $D, K, E, H$  are concyclic, just as we wished to prove.

---

<sup>1</sup>If you are not familiar with the power of a point, you can learn more in this handout (written in Czech): <https://prase.cz/library/MocnostboduokekruznicuAL/MocnostboduokekruznicuAL.pdf>.

POZNÁMKY:

Valná většina došlých řešení obdržela plný počet bodů. Z těchto úspěšných řešení valná většina nějakým způsobem využívala dokreslení bodu  $K$ , přičemž mnohá se namísto mocnosti k dokazované koncyklicitě dostala pouze úhlením. Při úhlení je často třeba dát si pozor na to, že zadání umožňuje více různých konfigurací bodů (v jakém pořadí leží na kružnici či na přímce), což vede k trochu odlišnému, ač v principu pořad stejnému úhlení. Opomenutí jedné nebo více takových konfigurací je jedním z nejspolehlivějších způsobů, jak např. v matematické olympiádě zbytečně ztratit body (ač v této úloze jsem za to nakonec body nestrhal). Způsobem, jak se rozebírání konfigurací vyvarovat, je *orientované úhlení*<sup>2</sup>. Lze si všimnout, že vzorové řešení tímto problémem netrpí, neboť lichoběžník má stejně dlouhá ramena, právě pokud má stejně dlouhé úhlopříčky (nezáleží na tom, jestli jsou  $DC$  a  $HK$  ramena, nebo úhlopříčky), a mocnost na konfiguraci nezávisí.

Jediným dalším funkčním dokreslením, které se v došlých řešeních vyskytlo, bylo využití bodu  $X$ , který je průsečíkem přímky  $DC$  s kružnicí opsanou  $DAB$  různým od  $D$ . (Matěj Doležálek)


**Problem 8.**

Let  $ABC$  be a non-equilateral triangle and  $G$  its centroid. Denote the midpoints of line segments  $AB$ ,  $CA$ ,  $BC$ ,  $AG$ ,  $CG$ , and  $BG$  by  $M_C$ ,  $M_B$ ,  $M_A$ ,  $N_A$ ,  $N_C$ , and  $N_B$  respectively. Show that the circumcircles of  $M_C N_A M_B$ ,  $M_B N_C M_A$ , and  $M_A N_B M_C$  all intersect in a single point.

(Radek Olšák)

SOLUTION:

Since  $M_A N_C$  is a midline of triangle  $BGC$ , it is parallel to  $BG$ . And as  $B$ ,  $N_B$ ,  $G$  and  $M_B$  lie on one line, it is also parallel to  $N_B M_B$ . Similarly,  $N_A M_C$  is parallel to  $N_B M_B$ , so we have  $N_A M_C \parallel M_A N_C$ . Analogously  $N_B M_A \parallel M_B N_A$  and  $N_C M_B \parallel M_C N_B$ .



Now we will use directed angles<sup>3</sup>. Let  $X$  be the intersection of circumcircles of  $M_B N_A M_C$  and  $M_B N_C M_A$  distinct from  $M_B$ . Then

$$\begin{aligned} \angle(M_A X, X M_C) &= \angle(M_A X, X M_B) + \angle(M_B X, X M_C) \\ &= \angle(M_A N_C, N_C M_B) + \angle(M_B N_A, N_A M_C) \\ &= \angle(N_B M_B, M_C N_B) + \angle(N_B M_A, M_B N_B) \\ &= \angle(M_A N_B, N_B M_C), \end{aligned}$$

<sup>2</sup>Viz např. <https://prase.cz/library/OrientovaneUhleniMO/OrientovaneUhleniMO.pdf>.

<sup>3</sup>Accessible from <https://prase.cz/library/OrientovaneUhlyMTa/OrientovaneUhlyMTa.pdf>.

where the second equality follows by concyclicity of  $M_B, N_A, M_C, X$  and  $M_B, N_C, M_A, X$ , and the third equality follows since  $M_A N_C \parallel N_B M_B$ ,  $N_A M_C \parallel N_B M_B$ ,  $N_B M_A \parallel M_B N_A$ , and  $N_C M_B \parallel M_C N_B$ . That means that  $M_A, N_B, M_C$ , and  $X$  are concyclic, which is what we wanted.

POZNÁMKY:

Většina přijatých řešení byla správně. Řešitelé si více či méně zdatně poradili s tím „jak vypadá obrázek“, přičemž nejjednodušší bylo postupovat přes orientované úhly, ale dalo se samozřejmě i jinak. (Rado van Švarc)